

Apendicitis en menores de cuatro años. Identificación de signos, síntomas y parámetros analíticos y radiológicos hacia un diagnóstico precoz

Publicado en Internet:
19-septiembre-2014

Ferran Campillo i López:
ferran.campillo@salud.madrid.org

E. García Camiño^a, F. Campillo i López^a, B. Delgado Díez^a,
E. Ballesteros Moya^a, A. Calle Gómez^a, J. Martín Sánchez^b

^aMIR-Pediatría. Hospital Universitario La Paz. Madrid, España

• ^bServicio de Urgencias Pediátricas. Hospital Universitario La Paz. Madrid, España.

Resumen

Introducción: la apendicitis aguda en menores de cuatro años presenta una incidencia elevada de complicaciones con respecto a otras edades, propias de un diagnóstico tardío, como la peritonitis. Con este estudio se pretende encontrar datos que ayuden a un diagnóstico precoz que reduzca su incidencia.

Material y métodos: estudio retrospectivo de los niños menores de cuatro años con diagnóstico de apendicitis aguda confirmada a partir de una muestra anatopatológica. Se registraron y analizaron variables demográficas, clínicas y analíticas, pruebas de imagen y anatopatológicas, y complicaciones quirúrgicas.

Resultados: de 82 pacientes incluidos, se encontró una relación de 1,5:1 hombre:mujer, que varió según la franja de edad, hallando un 85,7% de mujeres en menores de dos años y un 43,2% entre dos y cuatro años. Los síntomas más frecuentes fueron dolor abdominal progresivo, vómitos, anorexia y decaimiento. Los signos más frecuentes fueron dolor a la palpación en la fosa ilíaca derecha, defensa abdominal y fiebre. No hubo diferencias en cuanto a hallazgos analíticos, radiológicos o anatopatológicos.

La peritonitis fue más frecuente en los pacientes con fiebre >38,5 °C el primer día (*odds ratio*: 3,36; $p=0,009$). No se observó relación significativa entre edad y aumento de riesgo de peritonitis.

Conclusiones: la apendicitis aguda en menores de cuatro años sigue siendo un reto diagnóstico. Los síntomas y signos son similares a los clásicos de la apendicitis, aunque asocian más frecuentemente decaimiento y fiebre alta y de corta evolución; esta última triplica el riesgo de perforación apendicular y peritonitis, sobre todo en mujeres menores de dos años.

Palabras clave:

- Apendicitis
- Diagnóstico precoz
- Perforación intestinal
 - Peritonitis
 - Preescolar
 - Lactante

Appendicitis in children under four years. Identifying signs, symptoms, laboratory and imaging parameters for an early diagnosis

Abstract

Introduction: acute appendicitis in children under 4 years old shows a high incidence of complications due to a late diagnosis such as peritonitis, in comparison to other ages. This study is aimed to find data in order to achieve an early diagnosis to reduce their incidence.

Methodology: retrospective study for children under 4 years with confirmed diagnosis of acute appendicitis from pathological sample. Demographic, clinical, laboratory, imaging, anatomical pathology and surgical complications data were recorded and analyzed.

Results: eighty-two patients were enrolled, and a 1.5:1 ratio of male/female was found, which varied by age group, where 85.7% of female in children under 2 years and 43.2% between 2 and 4 years were found. The most frequent symptoms were progressive abdominal pain, vomiting, anorexia and malaise. The most common signs were tenderness in right lower quadrant, rebound pain and fever. There were no differences in laboratory, radiological or anatomical pathology findings.

 Texto íntegro en inglés disponible en: www.pap.es

Cómo citar este artículo: García Camiño E, Campillo i López F, Delgado Díez B, Ballesteros Moya E, Calle Gómez A, Martín Sánchez J. Apendicitis en menores de cuatro años. Identificación de signos, síntomas y parámetros analíticos y radiológicos hacia un diagnóstico precoz. Rev Pediatr Atención Primaria. 2014;16:213-8.

- Key words:**
- Appendicitis
 - Early diagnosis
 - Intestinal perforation
 - Peritonitis
 - Preschool child
 - Infant

Peritonitis was more frequent in patients with fever >38.5 °C on the first day (OR 3.36; p=0.009). No significant relationship was observed between age and increased risk of peritonitis.

Conclusions: acute appendicitis in children under 4 years is still a diagnostic challenge. Symptoms and signs are similar to classic appendicitis, though most frequently associated malaise and high fever and short evolution. Fever is related to a three times higher risk of appendical perforation and peritonitis, especially in women younger than 2 years.

INTRODUCCIÓN

El dolor abdominal agudo es un motivo de consulta frecuente en Pediatría de Atención Primaria, con un diagnóstico diferencial amplísimo que abarca desde patología banal a situaciones de riesgo vital. La apendicitis aguda en niños comparte características típicas de presentación con el adulto^{1,2}, a pesar de las diferencias anatómicas o dietéticas que modifican su fisiopatología³⁻⁵. Sin embargo, a medida que disminuye su edad se observan más frecuentemente complicaciones propias de un diagnóstico tardío, como la peritonitis secundaria a la perforación apendicular, especialmente en menores de cuatro años⁶⁻⁸.

Los objetivos del estudio fueron:

- Describir y analizar síntomas, signos y hallazgos analíticos, así como pruebas de imagen, en pacientes menores de cuatro años diagnosticados de apendicitis aguda.
- Identificar entre esos datos los más significativos para el diagnóstico precoz de apendicitis aguda.
- Encontrar entre esos datos los relacionados más frecuentemente con la principal complicación en estas edades: la perforación apendicular con peritonitis.
- Promover la recogida sistemática de los signos y síntomas que ayuden al clínico a conseguir un diagnóstico precoz.

MATERIAL Y MÉTODOS

Se realizó un estudio retrospectivo del periodo desde diciembre de 2006 a abril de 2012. Se incluyó a

menores de cuatro años atendidos en el Servicio de Urgencias Pediátricas de un hospital terciario universitario, con el diagnóstico de apendicitis aguda confirmado por estudio anatopatológico de la pieza quirúrgica. Los datos se obtuvieron de las historias a través de la Unidad de Codificación para pacientes con diagnóstico de apendicitis aguda al alta (CIE-10 K35.9) y edades inferiores a cuatro años en el momento del diagnóstico. Quedaron excluidas las apendicectomías realizadas por sospecha de apendicitis aguda que finalmente resultaron no patológicas (“apendicectomías blancas”), así como las profilácticas realizadas en el contexto de otra intervención. No se excluyó a aquellos niños que hubiesen sido diagnosticados en otros centros y que finalmente se trasladasen e interviniésem en el centro a estudio.

Se registraron variables demográficas, clínicas, analíticas, radiológicas, ecográficas, quirúrgicas y anatopatológicas, así como la estancia hospitalaria y las complicaciones posoperatorias, datos necesarios para la aplicación de alguna de las escalas validadas para el diagnóstico de apendicitis en niños, como el *Pediatric Appendicitis Score* (PAS) de M. Samuel⁹⁻¹¹.

En cuanto a la historia de la enfermedad, se reconocieron las horas de evolución del cuadro desde el primer día referido de síntomas hasta su diagnóstico, así como las visitas previas al hospital en ese periodo. Se consideraron los días de estancia desde el día de ingreso hasta el alta, ambos incluidos. El periodo de seguimiento de los pacientes abarcó desde la fecha del diagnóstico hasta la realización del estudio.

En relación a la radiología abdominal, se valoraron como hallazgos patológicos en la radiografía la dis-

tensión de asas, el aire libre intraabdominal, la presencia de apendicolito, el silencio abdominal y el borramiento de la línea del psoas; la escoliosis antiálgica fue calificada como no significativa. De la ecografía abdominal se estimó como alteración la presencia de apendicolito, absceso, plastrón y peritonitis (presencia de líquido purulento en la cavidad). También se valoraron como hallazgos anormales asas dilatadas, aumento del tamaño del apéndice, adenopatías inflamatorias, aumento de ecogenicidad de la grasa, apéndice no compresible y aumento del líquido en asas.

Respecto a las complicaciones, se consideraron tempranas aquellas ocurridas en los tres primeros días tras la cirugía, y tardías las posteriores a dicho periodo.

Finalmente, se llevó a cabo un análisis estadístico de las distintas variables de estudio con el programa Stata® 12.0.

RESULTADOS

En total se obtuvieron 82 pacientes que cumplían los criterios de elegibilidad. La edad media fue de 36,8 meses (rango 7-47 meses) y la mediana de 38,5 meses, y no se detectaron diferencias significativas entre las distintas variables según la edad. En nuestra muestra encontramos 48 varones y 34 mujeres, con una relación de 1,5:1. La relación varió según la franja de edad, encontrando un 85,7% de mujeres en los menores de dos años y un 43,2% entre los dos y los cuatro años.

La estancia media hospitalaria fue de nueve días. Para los menores de dos años, el diagnóstico se realizó de mediana a los dos días de evolución, y se demoró hasta el tercer día en el caso de los mayores. El 57,7% de los pacientes fue diagnosticado en su primera visita al hospital, el 30,7% en una segunda visita y hasta un 11,5% de los pacientes necesitó tres o más visitas para llegar al diagnóstico.

Los síntomas registrados con sus respectivas frecuencias se recogen en la **Tabla 1**. Los más frecuentes fueron dolor abdominal progresivo, vómitos, anorexia y decaimiento.

Tabla 1. Síntomas más frecuentes en menores de cuatro años con apendicitis aguda

Síntomas	Frecuencia (IC 95%)
Dolor abdominal:	
• Progresivo	96,2% (87,0 a 99,5)
• Intermitente	33,9% (20,8 a 47,1)
Vómitos	83,7% (73,8 a 91,0)
Anorexia	83% (68 a 93)
Decaimiento	70,7% (58,2 a 81,4)
Diarrea	37,3% (26,4 a 49,37)
Estreñimiento	8% (2,9 a 16,6)

IC 95%: intervalo de confianza del 95%.

Los signos más frecuentemente documentados fueron dolor a la palpación en el cuadrante inferior derecho o en la fosa iliaca derecha (FID), defensa abdominal y fiebre (**Tabla 2**). De este último síntoma cabe mencionar que el 74,1% de los pacientes que tenían fiebre alta ($>38,5^{\circ}\text{C}$) la desarrolló en el primer día de evolución.

En cuanto a los hallazgos de laboratorio (**Tabla 3**), destacaron proteína C reactiva (PCR) elevada, neutrófilia y leucocitosis. No hubo alteraciones en el contejo de plaquetas.

La ausencia de algunos de los datos en las historias clínicas, como la disuria o el dolor abdominal con las maniobras de Valsalva, impidió aplicar correctamente el PAS, por lo que no pudo ser incluido para su análisis.

En cuanto a las pruebas de imagen, se documentó que la radiografía simple de abdomen mostraba alteraciones hasta en el 75% de los pacientes, mientras que la ecografía abdominal fue patológica.

Tabla 2. Signos más frecuentes en menores de cuatro años con apendicitis aguda

Signos	Frecuencia (IC 95%)
Dolor a la palpación en la FID	84,5% (73,9 a 92,0)
Defensa abdominal	79,2% (68,4 a 87,6)
Fiebre	
• Desde el primer día	75,6% (64,6 a 84,6)
• $<38,5^{\circ}\text{C}$	24,4%
• $\geq38,5^{\circ}\text{C}$	51,3%
Dolor difuso a la palpación	72,6% (60,9 a 82,4)
Rebote positivo	64,4% (48,8 a 78,1)
Distensión abdominal	49,1% (35,1 a 63,2)

FID: fosa iliaca derecha; IC 95%: intervalo de confianza del 95%.

Tabla 3. Parámetros analíticos en menores de cuatro años con apendicitis aguda

Parámetros analíticos	Frecuencia (IC 95%)
Leucocitosis ($>15\,000/\mu\text{l}$)	60% (48,4 a 70,8)
Media	13 275 (12 000 a 14 500)
Neutrofilia ($>7000/\mu\text{l}$)	88,6% (79,5 a 94,7)
Plaquetas	Rangos normales 100%
PCR ($>5\,\text{mg/l}$)	97,4% (91,0 a 99,7)

IC 95%: intervalo de confianza del 95%; PCR: proteína C reactiva.

ca en un 88,8%. Los hallazgos ecográficos más frecuentemente encontrados fueron líquido libre/ peritonitis (26,9%) y signos inflamatorios (23,8%).

Los hallazgos anatomo-patológicos más frecuentes fueron apendicitis aguda gangrenosa (52,9%) y apendicitis aguda perforada (19,7%). Tanto en los mayores como en los menores de dos años, la anatomía patológica encontrada con mayor frecuencia fue la de apendicitis gangrenosa (71,4 y 62,2%, respectivamente).

Finalmente, se reportó peritonitis en un 58,4% de los pacientes (intervalo de confianza del 95% [IC 95%]: 46,6 a 69,5). Los pacientes con fiebre alta desde el primer día triplicaban el riesgo de desarrollar peritonitis (*odds ratio [OR]*: 3,3; IC 95%: 1,2 a 9,5; $p=0,009$). El resto de parámetros estudiados no presentó relación estadísticamente significativa con la peritonitis. Además, se documentaron abscesos abdominales en un 28,6% de los casos. No hubo ningún fallecimiento en la muestra estudiada.

DISCUSIÓN

Los datos obtenidos del estudio concuerdan en gran parte con otras series de características similares¹²⁻¹⁵. Sakellaris *et al.* estudiaron una muestra de 122 niños menores de cinco años con diagnóstico de apendicitis, que clasificaron en complicadas o no complicadas según la presencia de perforación apendicular en la intervención quirúrgica.

En ese estudio, el síntoma presente en prácticamente todos los casos fue el dolor abdominal de carácter difuso, junto con la presencia de fiebre, náuseas y vómitos, todos los cuales aparecieron

en, al menos, el 75% de los pacientes. Cerca de la mitad de los pacientes aquejaba anorexia y diarrea. Otros síntomas menos típicos fueron disuria, tos y odinofagia (como único síntoma en un 11%). A diferencia de nuestra serie, el decaimiento se encontró solo en el 6%. Los signos presentados más frecuentemente en las apendicitis complicadas fueron defensa abdominal (levemente superior en las apendicitis complicadas que en las no complicadas, del 76 y el 64%, respectivamente), signos de deshidratación (hasta en un 59%), distensión abdominal (30%), rebote (hasta en el 90% de las complicadas), ruidos hidroaéreos disminuidos y taquicardia. Otra circunstancia que influyó en el desarrollo de formas complicadas de apendicitis fue el retraso de la identificación de los síntomas más allá de la primera visita. Hasta un 66% de los pacientes que habían consultado en otra ocasión había sido diagnosticado de otro proceso, e incluso había recibido antibióticos.

Bundy *et al.* realizaron un metaanálisis de 42 estudios de apendicitis aguda en pacientes pediátricos de distintos rangos de edad. Tras el análisis, encontraron la fiebre como el mejor predictor para el diagnóstico de apendicitis en niños con dolor abdominal, hallazgo que coincide con el aumento significativo de hasta tres veces el riesgo para presentar peritonitis hallada en nuestra serie.

Un hallazgo llamativo de nuestra serie fue la predominancia del sexo femenino en los menores de dos años, que no se ha correspondido en otras muestras, y que puede deberse al pequeño tamaño muestral. El hecho de que el diagnóstico se realice antes en los menores de dos años con respecto a los mayores, puede deberse a la repercusión clínica de una fiebre alta de corta evolución acompañada de decaimiento en un paciente ya de por sí vulnerable, que lleva a la sospecha de una patología grave subyacente con posterior ingreso, aunque no se alcance el diagnóstico definitivo clínicamente.

En cuanto a las pruebas complementarias, el resto de las series coincide en la baja especificidad de los parámetros analíticos, aunque la ausencia de leucocitosis y neutrofilia hace el diagnóstico de apen-

dicitis poco probable. De igual manera, las pruebas de imagen no permiten el diagnóstico definitivo en todos los casos. No obstante, tanto la radiografía de abdomen simple como la ecografía abdominal se encuentran alteradas hasta en un 75% de los pacientes con apendicitis aguda.

Por lo que respecta al riesgo aumentado de desarrollar una apendicitis complicada en menores de cuatro años, son varios los estudios que se han fijado en la perforación del apéndice y posterior peritonitis^{3,12,16,17}. Kara *et al.* refieren hasta un 86% de perforaciones apendiculares en menores de un año, frecuencia que desciende hasta un 49% en niños de cuatro a cinco años. Se ha relacionado la formación de abscesos con la leucocitosis, la fiebre y la presencia de fecalito. Hasta el 66% de los niños menores de tres años de esa serie tenía perforación en el momento de la operación. Adicionalmente, la presentación de la enfermedad era más precoz, más grave y de mayor duración, y también se incrementaba la incidencia de complicaciones postquirúrgicas.

En las series revisadas, las complicaciones posoperatorias fueron poco frecuentes, destacando la formación de abscesos y las infecciones cutáneas. No se encontraron datos respecto a mortalidad.

La escasez de datos para la aplicación de la PAS evidencia la ausencia de estandarización en el manejo del dolor abdominal con sospecha de apendicitis, que no incluye sistemáticamente todos los ítems de las diferentes escalas validadas, además de las limitaciones propias de un estudio retrospectivo. De todas formas, las escalas validadas para niños en apendicitis no son específicas en niños tan pequeños.

La baja incidencia de esta entidad a estas edades y la inespecificidad de alguno de los síntomas parecen llevar al clínico a un bajo índice de sospecha y al retraso del diagnóstico, con la consiguiente demora en la intervención terapéutica.

CONCLUSIONES

El diagnóstico de apendicitis aguda en menores de cuatro años sigue siendo un reto diagnóstico. La elevada incidencia de perforaciones intestinales en este rango de edad, así como la formación de abscesos y otras complicaciones posoperatorias, estimula la búsqueda de herramientas para su diagnóstico precoz.

Los síntomas y signos son similares a los clásicos de la apendicitis, aunque podemos encontrar frecuentemente decaimiento y fiebre alta y de corta evolución en esta franja de edad, esta última triplica el riesgo de perforación apendicular y peritonitis, hallazgo más frecuente en las mujeres menores de dos años.

Los parámetros analíticos ayudan a descartar el diagnóstico, pero son poco específicos. De similar manera, las pruebas de imagen suelen estar alteradas, aunque pocas veces son concluyentes.

Finalmente, se hace evidente la necesidad de una recogida de datos sistemática y estandarizada de los parámetros usados en alguna de las escalas para el diagnóstico de la apendicitis aguda en niños, si bien no existen estudios que validen su uso en esta franja de edad concreta.

CONFLICTO DE INTERESES

Los autores declaran no presentar conflictos de intereses en relación con la preparación y publicación de este artículo.

Los resultados preliminares de este estudio se presentaron en la 18.^a Reunión de la Sociedad Española de Urgencias de Pediatría (SEUP), celebrada en Granada del 25 al 27 de abril de 2013, bajo el título “Diagnóstico de apendicitis aguda en menores de cuatro años”.

ABREVIATURAS

FID: fosa iliaca derecha • **IC 95%:** intervalo de confianza del 95% • **OR:** odds ratio • **PAS:** Pediatric Appendicitis Score • **PCR:** proteína C reactiva.

BIBLIOGRAFÍA

1. Wagner JM, McKinney WP, Carpenter JL. Does this patient have appendicitis? *JAMA*. 1996; 276:1589-94.
2. Mentine N, Guogenheim J. Appendicitis in the child and the adult. *La Revue du Praticien*. 2007;57:1947-52.
3. Jancelewicz T, Kim G, Miniati D. Neonatal appendicitis: a new look at an old zebra. *J Pediatr Surgery*. 2008;43:e1-5.
4. Barbosa AD, Júnior IF, Caetano RR, Lopes VG, Santos AM, Franco ED. Appendicitis in the premature newborn. *J Pediatr (Rio J)*. 2000;76:466-8.
5. Massad M, Srouji M, Awdeh A, Slim M, Tamer M, Tabbara M. Neonatal appendicitis: case report and a revised review of the English literature. *Z Kinderchir*. 1986;41:241-3.
6. Grosfeld JL, Weinberger M, Clatworthy HW. Acute appendicitis in the first two years of life. *J Pediatr Surgery*. 1973;8:285-93.
7. Schorlemmer GR, Herbst CA. Perforated neonatal appendicitis. *Southern Med J*. 1983;76:536-7.
8. Karaman A, CavuDođlu YH, Karaman I, Cakmak O. Seven cases of neonatal appendicitis with a review of the English language literature of the last century. *Pediatr Surgery Int*. 2003;19:707-9.
9. Samuel M. Pediatric appendicitis score. *J Pediatr Surgery*. 2002;37:877-81.
10. Goldman RD, Carter S, Stephens D, Antoon R, Mounstephen W, Langer JC. Prospective validation of the pediatric appendicitis score. *J Pediatr*. 2008;153:278-82.
11. Bhatt M, Joseph L, Ducharme FM, Dougherty G, McGillivray D. Prospective validation of the pediatric appendicitis score in a Canadian pediatric emergency department. *Acad Emerg Med*. 2009;16:591-6.
12. Sakellaris G, Tilemis S, Charassis G. Acute appendicitis in preschool-age children. *Eur J Pediatr*. 2005;164:80-3.
13. Bundy D, Byerley J, Liles E. Does this child have appendicitis? *JAMA*. 2007;298:438-51.
14. Mekhail P, Naguib N, Yanni F, Izzidien A. Appendicitis in paediatric age group: correlation between preoperative inflammatory markers and postoperative histological diagnosis. *Afr J Paediatr Surg*. 2011;8:309-12.
15. Brisighelli G, Morandi A, Parolini F, Leva E. Appendicitis in a 14-month-old infant with respiratory symptoms. *Afr J Paediatr Surg*. 2012;9: 148-51.
16. Hennelly KE, Bachur R. Appendicitis update. *Curr Opin Pediatr*. 2011;23:281-5.
17. Nelson DS, Bateman B, Bolte RG. Appendiceal perforation in children diagnosed in a pediatric emergency department. *Pediatr Emerg Care*. 2000;16:233-7.

Appendicitis in children under four years. Identifying signs, symptoms, laboratory and imaging parameters for an early diagnosis

E. García Camiño^a, F. Campillo i López^a, B. Delgado Díez^a,
E. Ballesteros Moya^a, A. Calle Gómez^a, J. Martín Sánchez^b

Published in Internet:
19-September-2014

Ferran Campillo i López:
ferran.campillo@salud.madrid.org

^aMedical Intern Resident (MIR), Paediatrics Department. Hospital Universitario La Paz. Madrid, Spain

^bPaediatric Emergency Department. Hospital Universitario La Paz. Madrid, Spain.

Abstract

Introduction: acute appendicitis in children under 4 years old shows a high incidence of complications due to a late diagnosis such as peritonitis, in comparison to other ages. This study is aimed to find data in order to achieve an early diagnosis to reduce their incidence.

Methodology: retrospective study for children under 4 years with confirmed diagnosis of acute appendicitis from pathological sample. Demographic, clinical, laboratory, imaging, anatomical pathology and surgical complications data were recorded and analyzed.

Results: eighty-two patients were enrolled, and a 1.5:1 ratio of male/female was found, which varied by age group, where 85.7% of female in children under 2 years and 43.2% between 2 and 4 years were found. The most frequent symptoms were progressive abdominal pain, vomiting, anorexia and malaise. The most common signs were tenderness in right lower quadrant, rebound pain and fever. There were no differences in laboratory, radiological or anatomical pathology findings.

Peritonitis was more frequent in patients with fever >38.5 °C on the first day (OR 3.36; p=0.009). No significant relationship was observed between age and increased risk of peritonitis.

Conclusions: acute appendicitis in children under 4 years is still a diagnostic challenge. Symptoms and signs are similar to classic appendicitis, though most frequently associated malaise and high fever and short evolution. Fever is related to a three times higher risk of appendical perforation and peritonitis, especially in women younger than 2 years.

- Key words:**
- Appendicitis
 - Early diagnosis
 - Intestinal perforation
 - Peritonitis
 - Preschool
 - Infant

Apendicitis en menores de cuatro años. Identificación de signos, síntomas y parámetros analíticos y radiológicos hacia un diagnóstico precoz

Resumen

Introducción: la apendicitis aguda en menores de cuatro años presenta una incidencia elevada de complicaciones con respecto a otras edades, propias de un diagnóstico tardío, como la peritonitis. Con este estudio se pretende encontrar datos que ayuden a un diagnóstico precoz que reduzca su incidencia.

Material y métodos: estudio retrospectivo de los niños menores de cuatro años con diagnóstico de apendicitis aguda confirmada a partir de una muestra anatopatológica. Se registraron y analizaron variables demográficas, clínicas y analíticas, pruebas de imagen y anatopatológicas, y complicaciones quirúrgicas.

Resultados: de 82 pacientes incluidos, se encontró una relación de 1,5:1 hombre:mujer, que varió según la franja de edad, hallando un 85,7% de mujeres en menores de dos años y un 43,2% entre dos y cuatro años. Los síntomas más frecuentes fueron dolor abdominal progresivo, vómitos, anorexia y decaimiento. Los signos más frecuentes fueron dolor a la palpación en la fosa ilíaca derecha, defensa abdominal y fiebre. No hubo diferencias en cuanto a hallazgos analíticos, radiológicos o anatopatológicos.

How to quote this article: García Camiño E, Campillo López F, Delgado Díez B, Ballesteros Moya E, Calle Gómez A, Martín Sánchez J. Appendicitis in children under four years. Identifying signs, symptoms, laboratory and imaging parameters for an early diagnosis. Rev Pediatr Aten Primaria. 2014;16:213-8.

- Palabras clave:**
- Apendicitis
 - Diagnóstico precoz
 - Perforación intestinal
 - Peritonitis
 - Preescolar. Lactante

La peritonitis fue más frecuente en los pacientes con fiebre >38,5 °C el primer día (*odds ratio*: 3,36; $p=0,009$). No se observó relación significativa entre edad y aumento de riesgo de peritonitis.

Conclusiones: la apendicitis aguda en menores de cuatro años sigue siendo un reto diagnóstico. Los síntomas y signos son similares a los clásicos de la apendicitis, aunque asocian más frecuentemente decaimiento y fiebre alta y de corta evolución; esta última triplica el riesgo de perforación apendicular y peritonitis, sobre todo en mujeres menores de dos años.

INTRODUCTION

Acute abdominal pain is a frequent reason for visits to primary care paediatrics departments, and has an extensive differential diagnosis that ranges from common mild pathologies to life-threatening conditions. Some of the features of acute appendicitis in children are also typical of adult presentations,^{1,2} despite the anatomical or dietary differences that make the pathophysiology of these two groups different.³⁻⁵ However, the younger the patient, the more common it is to observe complications characteristic of a delayed diagnosis, such as peritonitis secondary to perforated appendicitis, especially in children younger than 4 years.⁶⁻⁸

The objectives of the study were:

- Describing and analysing symptoms, signs, laboratory findings, and imaging tests in patients younger than 4 years diagnosed with acute appendicitis.
- Identifying the data that are most significant for the early diagnosis of acute appendicitis.
- Identifying the data most frequently associated with the main complication in this age group: appendiceal perforation with peritonitis.
- Promoting the systematic documentation of signs and symptoms that can help the clinician make an early diagnosis.

MATERIALS AND METHODS

We conducted a retrospective study that ran from December 2006 to April 2012. The study included

children younger than 4 years seen at the paediatric emergency department of a tertiary university hospital with a diagnosis of acute appendicitis confirmed by histological examination of the appendix after surgery. We obtained the data from the medical records of patients with a discharge diagnosis of acute appendicitis (ICD-10 K35.9) from the hospital's coding unit. We excluded cases in which appendectomy was performed for suspected acute appendicitis if they turned out to be nonpathological ("white appendectomies"), as well as cases in which prophylactic appendectomies were performed in the context of another intervention. The study did not exclude patients that were diagnosed in other centres and eventually transferred to our hospital for surgery.

We collected data for demographic, clinical, laboratory, radiological, ultrasonographic, and anatomical pathology variables, and on the length of stay and postoperative complications, data needed to apply some of the validated scales for diagnosing appendicitis in children, such as the Pediatric Appendicitis Score (PAS) developed by M. Samuel.⁹⁻¹¹

When it came to the evolution of the disease, we collected the duration of symptoms from the reported day of onset to the day of diagnosis, as well as any visits to the hospital that occurred in this time interval. We collected the length of stay in days from admission to discharge, including both. The follow-up period comprehended the time from the day of diagnosis to the time of the study.

The data collected for the abnormal findings of abdominal radiographs included presence of dilated bowel loops, free intraabdominal air, appendicolith, silent abdomen, and obliteration of the psoas

shadow; we did not consider antalgic scoliosis significant. Findings considered abnormal in the abdominal ultrasound were presence of appendicolith, abscess, appendiceal mass and peritonitis (presence of suppurative fluid in the peritoneal cavity). We also analysed abnormal findings such as dilated bowel loops, enlarged appendix, inflammatory adenopathies, increased echogenicity of periappendiceal fat, noncompressible appendix, and fluid collection in bowel loops.

We classified complications as early if they occurred in the first three days postsurgery, and late if they developed afterward.

Lastly, we analysed the different study variables using the Stata® software, version 12.0.

RESULTS

The study included a total of 82 patients that met the eligibility criteria. The mean age was 36.8 months (range, 7–47 months) with a median of 38.5 months. We found no significant differences in the various variables between age groups. The sample included 48 boys and 34 girls, resulting in a 1.5:1 ratio. This ratio varied depending on age, with girls accounting for 85.7% of patients under 2 years of age and 43.2% of patients between 2 and 4 years of age.

The mean length of stay was 9 days. The median time to diagnosis was 2 days since the onset of symptoms in children younger than 2 years, and extended to 3 days for older children. Of all patients, 57.7% were diagnosed in their first visit to the hospital, 30.7% during a second visit, and up to 11.5% required three or more visits to get diagnosed.

Table 1 presents the documented symptoms and their respective frequencies. The most frequent symptoms were progressive abdominal pain, vomiting, anorexia, and asthenia.

The most frequent signs were pain upon palpation of the lower right quadrant or the right iliac fossa (RIF), abdominal guarding, and fever (**Table 2**). In regards to the latter, we ought to note that 74.1%

Table 1. Most frequent symptoms in children younger than 4 years with acute appendicitis

Symptoms	Frequency (95% CI)
Abdominal pain:	
• Progressive	96.2% (87.0–99.5)
• Intermittent	33.9% (20.8–47.1)
Vomiting	83.7% (73.8–91.0)
Anorexia	83% (68–93)
Asthenia	70.7% (58.2–81.4)
Diarrhoea	37.3% (26.4–49.37)
Constipation	8% (2.9–16.6)

95% CI: 95% confidence interval.

of patients presenting with high fever (>38.5 °C) had developed it on the day of the onset of symptoms.

The main laboratory findings (**Table 3**) were elevated C-reactive protein (CRP), neutrophilia and leukocytosis. The platelet counts were normal.

The absence of some data in the medical records, such as dysuria or abdominal pain upon performing the Valsalva manoeuvre, precluded the correct application of the PAS, so we could not include this score in our analysis.

When it came to imaging tests, the documentation showed that plain radiography of the abdomen showed abnormalities in up to 75% of patients, while the abdominal ultrasound was abnormal in 88.8%. The most frequent ultrasound findings were free fluid/peritonitis (26.9%) and signs of inflammation (23.8%).

The most common anatomical pathology findings were acute gangrenous appendicitis (52.9%) and

Table 2. Most frequent signs in children younger than 4 years with acute appendicitis

Signs	Frequency (95% CI)
Pain upon palpation of the RIF	84.5% (73.9–92.0)
Abdominal guarding	79.2% (68.4–87.6)
Fever	
• Starting on day of onset	75.6% (64.6 a 84.6)
• <38.5 °C	24.4%
• ≥38.5 °C	51.3%
Diffuse pain upon palpation	72.6% (60.9–82.4)
Rebound tenderness	64.4% (48.8–78.1)
Abdominal distension	49.1% (35.1–63.2)

RIF: right iliac fossa; 95% CI: 95% confidence interval.

Table 3. Laboratory parameters in children younger than 4 years with acute appendicitis

Laboratory parameters	Frequency (95% CI)
Leukocytosis (>15 000/ μ l)	60% (48.4–70.8)
Mean count	13 275 (12 000–14 500)
Neutrophilia (>7000/ μ l)	88.6% (79.5–94.7)
Platelets	Normal ranges 100%
CRP (>5 mg/l)	97.4% (91.0–99.7)

CRP: C-reactive protein; 95% CI: 95% confidence interval.

acute perforated appendicitis (19.7%). Gangrenous appendicitis was the most common finding in anatomical pathology in both children older and younger than two years (71.4% and 62.2%, respectively).

Lastly, peritonitis was documented in 58.4% of the patients (95% confidence interval [95% CI]: 46.6–69.5). Patients who developed a high fever on the day of onset had three times the risk of developing peritonitis (odds ratio [OR]: 3.3; 95% CI: 1.2–9.5; $p=0.009$). The rest of the parameters under study showed no statistically significant relationship with peritonitis. Also, abdominal abscesses were documented in 28.6% of the cases. There were no deaths in the sample under study.

DISCUSSION

The data obtained in our study are mostly consistent with those found in other case series of similar characteristics.^{12–15} Sakellaris *et al* studied a sample of 122 children younger than 5 years diagnosed with appendicitis, which they classified as complicated or uncomplicated depending on the presence of appendiceal perforation at surgery. In their study, the symptom that was present in almost all cases was diffuse abdominal pain accompanied by fever, nausea and vomiting, all of which developed in at least 75% of the patients. Nearly half of the patients presented with anorexia and diarrhoea. Less typical symptoms included dysuria, cough, and odynophagia (as the only symptom in 11%). Unlike what happened in our series, asthenia was only observed in 6%. The signs found most frequently in complicated appendicitis cases were

abdominal guarding (slightly higher in complicated versus uncomplicated appendicitis, in 76% and 64% of cases, respectively), signs of dehydration (in up to 59%), abdominal distension (30%), rebound tenderness (in up to 90% of complex cases), decreased bowel sounds and tachycardia. Another circumstance that contributed to the development of complicated forms of appendicitis was the delayed identification of symptoms beyond the first visit. Up to 66% of the patients that had sought care at a prior time had been diagnosed with a different condition, and had even been treated with antibiotics.

Bundy *et al* performed a literature review of 42 studies on acute appendicitis in children of different age ranges. Their analysis showed that fever was the best diagnostic predictor for acute appendicitis in children with abdominal pain, a finding that is consistent with the significant increase of up to three times in the risk for peritonitis found in our series.

An unexpected finding in our study was the predominance of the female sex in patients younger than 2 years that has not been found in other samples, which may have been due to the small sample size. The fact that children younger than two years are diagnosed earlier than the older children in our sample may be due to the clinical importance of a high fever of rapid onset accompanied by asthenia in these patients that are vulnerable in themselves, which leads to suspicion of a severe underlying pathology and admission to the hospital, even if the clinical signs and symptoms are not sufficient to make a definitive diagnosis.

As for diagnostic tests, previous case series are consistent in showing low specificity for the laboratory parameters, although the absence of leukocytosis and neutrophilia makes the diagnosis of appendicitis improbable. Likewise, imaging tests do not allow a definitive diagnosis in every case. Nevertheless, both plain abdominal radiographs and abdominal ultrasound scans are abnormal in up to 75% of patients with acute appendicitis.

On the subject of the increased risk of developing complicated appendicitis of children younger than

4 years, several studies have focused on appendiceal perforation and subsequent peritonitis.^{3,12,16,17} Kara *et al* reported the development of appendiceal perforation in up to 86% of children younger than 1 year, a frequency that decreased to as low as 49% in children aged 4 and 5 years. Abscess formation was associated with leukocytosis, fever, and the presence of faecalith. Up to 66% of children younger than 3 years in this series had perforation at the time of surgery. Furthermore, their symptoms developed earlier, were more severe, and lasted longer, and the incidence of postoperative complications was also higher in this group.

Postoperative complications were infrequent in all the case series we reviewed, and the most common ones were abscess formation and skin infections. We did not find data on mortality.

The scarcity of the data required for the application of the PAS evinces the lack of standardisation in the management of abdominal pain in cases of suspected appendicitis, as the management does not systematically gather information for all the items in the different validated scales, in addition to the limitations characteristic of a retrospective study. At any rate, the scales validated for appendicitis in children are not specific in children this young.

The low incidence of this entity in this age group and the nonspecificity of some of its symptoms result in a low level of suspicion on the part of the clinician and in delayed diagnosis, with the consequent delay in therapeutic intervention.

CONCLUSIONS

The diagnosis of acute appendicitis in children younger than 4 years continues to be a challenge.

The high incidence of bowel perforation in this age group, as well as the development of abscesses and other postoperative complications, motivates the search for tools to aid in its early diagnosis.

The signs and symptoms in this age group are similar to the classic signs and symptoms of appendicitis, although asthenia and high fever of short duration is found often in this group, as well as three times the risk of bowel perforation and peritonitis, a finding that is more frequent in girls younger than two years.

Laboratory parameters help rule out the diagnosis, but their specificity is low. Similarly, imaging studies are usually abnormal, but seldom conclusive.

Finally, it is necessary that data for the parameters used in some of the scales for the diagnosis of acute appendicitis in children be collected in a systematic and standardised manner, even though there are no studies yet validating their use in this particular age group.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare in relation to the preparation and publication of this paper.

Preliminary results of this study were presented at the 18 Reunión de la Sociedad Española de Urgencias de Pediatría (SEUP), held in Granada from 25 to 27 de April 2013, under the title "Diagnóstico de apendicitis aguda en menores de cuatro años".

ACRONYMS

95% CI: 95% confidence interval • CRP: C-reactive protein
 • OR: odds ratio • PAS: Pediatric Appendicitis Score • RIF: right iliac fossa.

BIBLIOGRAPHY

1. Wagner JM, McKinney WP, Carpenter JL. Does this patient have appendicitis? *JAMA*. 1996;276:1589-94.
2. Mentine N, Guogenheim J. Appendicitis in the child and the adult. *La Revue du Praticien*. 2007;57:1947-52.
3. Jancelewicz T, Kim G, Miniati D. Neonatal appendicitis: a new look at an old zebra. *J Pediatr Surgery*. 2008;43:e1-5.
4. Barbosa AD, Júnior IF, Caetano RR, Lopes VG, Santos AM, Franco ED. Appendicitis in the premature newborn. *Jornal de pediatria*. 2000;76:466-8.
5. Massad M, Srouji M, Awdeh A, Slim M, Tamer M, Tabbara M. Neonatal appendicitis: case report and a revised review of the English literature. *Z Kinderchir*. 1986;41:241-3.
6. Grosfeld JL, Weinberger M, Clatworthy HW. Acute appendicitis in the first two years of life. *J Pediatr Surgery*. 1973;8:285-93.
7. Schorlemmer GR, Herbst CA. Perforated neonatal appendicitis. *Southern Med J*. 1983;76:536-7.
8. Karaman A, CavuDođlu YH, Karaman I, Cakmak O. Seven cases of neonatal appendicitis with a review of the English language literature of the last century. *Pediatr Surgery Int*. 2003;19:707-9.
9. Samuel M. Pediatric appendicitis score. *J Pediatr Surgery*. 2002;37:877-81.
10. Goldman RD, Carter S, Stephens D, Antoon R, Mounstephen W, Langer JC. Prospective validation of the pediatric appendicitis score. *J Pediatr*. 2008;153:278-82.
11. Bhatt M, Joseph L, Ducharme FM, Dougherty G, McGillivray D. Prospective validation of the pediatric appendicitis score in a Canadian pediatric emergency department. *Acad Emerg Med*. 2009;16:591-6.
12. Sakellaris G, Tilemis S, Charassis G. Acute appendicitis in preschool-age children. *Eur J Pediatr*. 2005;164:80-3.
13. Bundy D, Byerley J, Liles E. Does this child have appendicitis? *JAMA*. 2007;298:438-51.
14. Mekhail P, Naguib N, Yanni F, Izzidien A. Appendicitis in paediatric age group: correlation between preoperative inflammatory markers and postoperative histological diagnosis. *Afr J Paediatr Surg*. 2011;8:309-12.
15. Brisighelli G, Morandi A, Parolini F, Leva E. Appendicitis in a 14-month-old infant with respiratory symptoms. *Afr J Paediatr Surg*. 2012;9:148-51.
16. Hennelly KE, Bachur R. Appendicitis update. *Curr Opin Pediatr*. 2011;23:281-5.
17. Nelson DS, Bateman B, Bolte RG. Appendiceal perforation in children diagnosed in a pediatric emergency department. *Pediatr Emerg Care*. 2000;16:233-7.